

## **Cardiac reserve, Cardiac work & Oxygen consumption by the heart**

### **ILOs**

By the end of this lecture the students will be able to:

1. By the end of this lecture the students will be able to:
2. Describe cardiac reserve with its parameters limitations, and mechanisms
3. Describe cardiac output curve , hypo-effective and hyper-effective hearts and their causes.
4. List the types of work done by the heart
5. Describe mechanical efficiency of the heart
6. Describe the O<sub>2</sub> consumption by the heart
7. Apply the information studied in this section to solve a clinical problem or explain clinical case.

### **Cardiac Reserve**

- ✓ It is the ability of the heart to augment its CO in order to meet the body demands.
- ✓ It equal the difference between basal and maximal cardiac output or work.
- ✓ Cardiac reserve = Maximal cardiac output - basal cardiac output  
= 35 (L/min) - 5 (L/min).  
= 30 (L/min).

### **N.B:**

- ✓ As the venous return increase to the heart, cardiac output increases up to a limit after which the heart will become a limiting factor for increased cardiac output.
- ✓ Normal cardiac output curve show that heart can increase cardiac output up to 13 L/min ( that is 2.5 times the normal cardiac output) before plateau is obtained.
- ✓ This mean that to increase cardiac output above 13 L/min, specific way of heart stimulation are needed.

### **Mechanisms of cardiac reserve :**

This happen through increasing the determinants of CO: stroke volume and heart rate.

## **A) Short acting mechanisms:**

Moment to moment increase in CO to meet the increased demands.

### **1) Increase CO up to 13 L/min (permissive limit):**

#### **a) Heart rate increase:**

- ✓ from basal 60 beat up to 90 beats which is the intrinsic pacemaker frequency.
- ✓ Happen by inhibition of vagal tone without sympathetic stimulation.

#### **b) Stroke volume:**

Increased by increased venous return and EDV by heterometric mechanism (Frank-Starling law). Followed by homometric regulation (with same EDV) if condition prolonged than 2 minutes, SV is increased by decreased ESV.

### **2) Increase CO above the permissive limit:**

This depends on sympathetic stimulation and circulating catecholamines. It is due to  $\beta_1$  receptor stimulation.

#### **a) Heart rate reserve mechanism:**

Sympathoadrenal stimulation exerts a +ve chronotropic effect and heart rate can increase up to 180 beat per minute.

#### **b) Stroke volume reserve mechanism:**

Sympathoadrenal stimulation exerts a +ve inotropic effect with constant EDV, SV increases by increasing power of contraction and decreasing ESV down to 30 ml.

## **B) Long term Mechanisms :**

### **Slowly and gradual acting mechanism:**

#### **1) Dilation :**

- ✓ Depend on Frank-Starling law increasing EDV.
- ✓ Occur in volume overload as aortic incompetence, ventricular septal defect and congestive heart failure.

#### **2) Hypertrophy:**

- ✓ Increased number and thickness of cardiac muscles, increasing the thickness of ventricular wall to overcome the excessive resistance.
- ✓ Occur in pressure overload as in hypertension, aortic stenosis.

### **Limitation of cardiac reserve:**


- ✓ Increased CO up to permissive limit (13L/min) has no limitation.
- ✓ Sympathoadrenal stimulation is limited by stores of norepinephrine and responsiveness to  $\beta$  adrenergic receptors.

- ✓ If heart rate increased above 180 beat per minute:
  - ✓ cardiac output is decreased due to reduced ventricular filling and consequently stroke volume.
  - ✓ Increased work of the heart and O<sub>2</sub> consumption while coronary blood flow is decreased.
- ✓ If EDV is excessively increased, overstretch of the wall with extra need for oxygen occur leading to diastolic dysfunctions.
- ✓ If contractility is excessively increased, injury of myocardium may happen as in athlete heart.
- ✓ If hypertrophy exceeded certain limit, decreased blood supply occur as blood vessels doesn't increase in the same extent as muscle hypertrophy occur (disproportionate).

### **Cardiac output curve:**

- ✓ The normal cardiac output curve, showing the cardiac output per minute at each level of right atrial pressure.
- ✓ Causes of hyper-effective heart:  
Hyper-effective heart is heart with increased cardiac output better than normal. It can be caused by:
  - 1) Nervous stimulation.
    - Nervous excitation through parasympathetic inhibition and sympathetic stimulation, increase heart rate and power of contraction. This can increase cardiac output up to 25 L/min.
  - 2) Hypertrophy.
    - Work over load for the heart during exercise cause cardiac hypertrophy.
    - In marathon runners, hypertrophy occurs by 50- 75% which leads to increase cardiac output up to 60-100%.

In marathon runners, both factors are combined so CO can increase up to 30 - 40 L/min.
- ✓ Causes of hypo-effective heart:  
Hypo-effective heart is heart with decreased cardiac output. It can be caused by:
  - 1) Increased arterial pressure such as in severe hypertension.
  - 2) Inhibition of nervous excitation of the heart.
  - 3) Arrhythmias.
  - 4) Coronary artery insufficiency.
  - 5) Valvular heart disease.
  - 6) Congenital heart disease.
  - 7) Myocarditis.
  - 8) Cardiac hypoxia.



### Cardiac work

- ✓ Work is the product of force times distance.
- ✓ In the case of the work done to move a volume of fluid, work is defined as the product of the volume of fluid and the pressure required to move the fluid.

#### Types of cardiac work:

##### ✓ Stroke Work:


- is the product of stroke volume and pressure needed to eject it (equal to mean systolic pressure during ejection).
- **Stroke work = stroke volume × mean arterial pressure.**
- Left ventricle have stroke work 7 times than that of right ventricle as aortic pressure is 7 times the pulmonary pressure.

##### ✓ Minute work:

- is work done by heart in whole minute.
- Minute work = stroke work × heart rate.

- ✓ Increasing the pressure with constant stroke volume increase the cardiac work (pressure work).
- ✓ Increasing the cardiac output (stroke volume) with constant pressure increase the cardiac work (volume work).
- ✓ Pressure work consume more O<sub>2</sub> than volume work, so the heart with coronary insufficiency can have angina pectoris more common with aortic stenosis (pressure load) more than with aortic incompetence ( volume load).
- ✓ So, work output of heart is in form of
  - External work : include pressure and volume work.
  - Kinetic energy to give velocity to the blood ( normally small and neglected = 1%. But increased in case of aortic stenosis).

- And heat.



Guyton and Hall 13<sup>th</sup> edition

### **Mechanical efficiency of the heart :**


- ✓ The efficiency of the heart may be calculated as the ratio of the work accomplished to the total energy used. Normally, it is 20-25%.
- ✓ It is decreased with increased after load, preload, heart rate and heart dilation.
- ✓ It is increased with hypertrophy.

### **Oxygen consumption by the heart**

- ✓ O<sub>2</sub> consumption by the beating heart is about 9 mL/100 g/min at rest.
- ✓ O<sub>2</sub> consumption increases in exercise and in a number of different states.
- ✓ O<sub>2</sub> extraction by cardiac muscles is high and little additional O<sub>2</sub> can be extracted from the blood in the coronaries. so increases in O<sub>2</sub> consumption require increases in coronary blood flow.

### Factors affecting O<sub>2</sub> consumption:

- ✓ O<sub>2</sub> consumption by the heart is determined primarily by the intramyocardial tension, the contractile state of the myocardium, and the heart rate.
- ✓ When the heart rate is increased by sympathetic stimulation, O<sub>2</sub> consumption is increased due to increased number of beats, increased velocity and strength of each contraction.
- ✓ When heart is dilated, according to Laplace law, tension developed in the wall is increased and accordingly O<sub>2</sub> consumption is increased.
- ✓ O<sub>2</sub> consumption is increased with increase in after load and increased preload. But Afterload produce more O<sub>2</sub> consumption.



Guyton and Hall 13<sup>th</sup> edition